

Sharp two parameter bounds for logarithmic and arithmetic-geometric means

Yu-Ming Chu¹, Ye-Fang Qiu¹, Miao-Kun Wang¹ and Xiao-Yan Ma²

¹Department of Mathematics, Huzhou Teachers College, Huzhou 313000, China;

²Department of Mathematics, Zhejiang Sci-Tech University, Hangzhou 310018, China.

Correspondence should be addressed to Yu-Ming Chu, chuyuming@hutczj.cn

Abstract: For fixed $s \geq 1$ and $t_1, t_2 \in (0, 1/2)$ we prove that the inequalities $G^s(t_1a + (1 - t_1)b, t_1b + (1 - t_1)a)A^{1-s}(a, b) > AG(a, b)$ and $G^s(t_2a + (1 - t_2)b, t_2b + (1 - t_2)a)A^{1-s}(a, b) > L(a, b)$ hold for all $a, b > 0$ with $a \neq b$ if and only if $t_1 \geq 1/2 - \sqrt{2s}/(4s)$ and $t_2 \geq 1/2 - \sqrt{6s}/(6s)$. Here $G(a, b)$, $L(a, b)$, $AG(a, b)$ and $A(a, b)$ are the geometric, logarithmic, arithmetic-geometric and arithmetic means of a and b , respectively.

2010 Mathematics Subject Classification: 26E60.

Keywords: geometric mean, logarithmic mean, arithmetic-geometric mean, arithmetic mean.

1 Introduction

For real numbers a, b and c with $c \neq 0, -1, -2, \dots$, the Gaussian hypergeometric function is defined by

$$F(a, b; c; x) = {}_2F_1(a, b; c; x) = \sum_{n=0}^{\infty} \frac{(a, n)(b, n)}{(c, n)} \frac{x^n}{n!}, \quad |x| < 1. \quad (1.1)$$

Here $(a, 0) = 1$ for $a \neq 0$, and $(a, n) = a(a+1)(a+2)(a+3) \cdots (a+n-1)$ is the shifted factorial function for $n = 1, 2, \dots$. In connection with the Gaussian hypergeometric function, the well-known complete elliptic integrals $\mathcal{K}(r)$ and $\mathcal{E}(r)$ ($0 < r < 1$) of the first and second kinds [1, 2] are defined by

$$\begin{cases} \mathcal{K}(r) = \pi F(1/2, 1/2; 1; r^2)/2 = \int_0^{\pi/2} (1 - r^2 \sin^2 \theta)^{-1/2} d\theta, \\ \mathcal{K}(0) = \pi/2, \quad \mathcal{K}(1) = \infty \end{cases} \quad (1.2)$$

and

$$\begin{cases} \mathcal{E}(r) = \pi F(-1/2, 1/2; 1; r^2)/2 = \int_0^{\pi/2} (1 - r^2 \sin^2 \theta)^{1/2} d\theta, \\ \mathcal{E}(0) = \pi/2, \quad \mathcal{E}(1) = 1, \end{cases} \quad (1.3)$$

respectively. The following formulas for $\mathcal{K}(r)$ were presented in [3]:

$$\frac{d\mathcal{K}(r)}{dr} = \frac{\mathcal{E}(r) - (1 - r^2)\mathcal{K}(r)}{r(1 - r^2)}, \quad (1.4)$$

$$\mathcal{K}\left(\frac{2\sqrt{r}}{1+r}\right) = (1+r)\mathcal{K}(r). \quad (1.5)$$

Let $H(a, b) = 2ab/(a+b)$, $G(a, b) = \sqrt{ab}$, $L(a, b) = (b-a)/(\log a - \log b)$ and $A(a, b) = (a+b)/2$ be the classical harmonic, geometric, logarithmic and arithmetic means of two distinct positive real numbers a and b , respectively. Then it is well known that the inequalities $H(a, b) < G(a, b) < L(a, b) < A(a, b)$ hold for all $a, b > 0$ with $a \neq b$.

The classical arithmetic-geometric mean $AG(a, b)$ of two positive number a and b is defined as the common limit of sequences $\{a_n\}$ and $\{b_n\}$, which are given by

$$\begin{aligned} a_0 &= a, & b_0 &= b, \\ a_{n+1} &= (a_n + b_n)/2 = A(a_n, b_n), & b_{n+1} &= \sqrt{a_n b_n} = G(a_n, b_n). \end{aligned}$$

It is well known that inequalities

$$G(a, b) < \sqrt{A(a, b)G(a, b)} < AG(a, b) < A(a, b) \quad (1.6)$$

hold for all $a, b > 0$ with $a \neq b$.

Recently, the harmonic, geometric, logarithmic, arithmetic-geometric and arithmetic means have been the subject of intensive research. In particular, many remarkable inequalities for these means can be found in the literature [4-13].

The Gaussian identity [3] shows that

$$AG(1, r)\mathcal{K}(\sqrt{1-r^2}) = \frac{\pi}{2} \quad (1.7)$$

for all $r \in (0, 1)$.

Carlson and Vuorinen [14], and Brackenn [15] proved that

$$L(a, b) < AG(a, b)$$

for all $a, b > 0$ with $a \neq b$. Vamanamurthy and Vuorinen [16] established that

$$AG(a, b) < \frac{\pi}{2} L(a, b)$$

for all $a, b > 0$ with $a \neq b$.

For $t_1, t_2, t_3, t_4 \in (0, 1/2)$, very recently Chu et al. [17, 18] proved that the inequalities

$$G(t_1 a + (1 - t_1)b, t_1 b + (1 - t_1)a) > AG(a, b), \quad (1.8)$$

$$H(t_2 a + (1 - t_2)b, t_2 b + (1 - t_2)a) > AG(a, b), \quad (1.9)$$

$$G(t_3 a + (1 - t_3)b, t_3 b + (1 - t_3)a) > L(a, b), \quad (1.10)$$

and

$$H(t_4 a + (1 - t_4)b, t_4 b + (1 - t_4)a) > L(a, b) \quad (1.11)$$

hold for all $a, b > 0$ with $a \neq b$ if and only if $t_1 \geq 1/2 - \sqrt{2}/4$, $t_2 \geq 1/4$, $t_3 \geq 1/2 - \sqrt{6}/6$ and $t_4 \geq 1/2 - \sqrt{3}/6$.

Let $t \in (0, 1/2)$, $s \geq 1$ and

$$Q_{t,s}(a, b) = G^s(ta + (1 - t)b, tb + (1 - t)a) A^{1-s}(a, b). \quad (1.12)$$

Then it is not difficult to verify that

$$Q_{t,1}(a, b) = G(ta + (1 - t)b, tb + (1 - t)a),$$

$$Q_{t,2}(a, b) = H(ta + (1 - t)b, tb + (1 - t)a)$$

and $Q_{t,s}(a, b)$ is strictly increasing with respect to $t \in (0, 1/2)$ for fixed $a, b > 0$ with $a \neq b$.

It is natural to ask what are the least values $t_1 = t_1(s)$ and $t_2 = t_2(s)$ in $(0, 1/2)$ such that inequalities $Q_{t_1,s}(a, b) > AG(a, b)$ and $Q_{t_2,s}(a, b) > L(a, b)$ hold for all $a, b > 0$ with $a \neq b$ and $s \geq 1$. The aim of this paper is to answer these questions, our main results are the following Theorems 1.1 and 1.2.

Theorem 1.1. If $t \in (0, 1/2)$ and $s \geq 1$ then the inequality

$$Q_{t,s}(a, b) > AG(a, b) \quad (1.13)$$

holds for all $a, b > 0$ with $a \neq b$ if and only if $t \geq 1/2 - \sqrt{2s}/(4s)$.

Theorem 1.2. If $t \in (0, 1/2)$ and $s \geq 1$ then the inequality

$$Q_{t,s}(a, b) > L(a, b) \quad (1.14)$$

holds for all $a, b > 0$ with $a \neq b$ if and only if $t \geq 1/2 - \sqrt{6s}/(6s)$.

Remark 1.1. Let $s = 1, 2$ in Theorem 1.1, then inequality (1.13) becomes inequalities (1.8) and (1.9), respectively.

Remark 1.2. Let $s = 1, 2$ in Theorem 1.2, then inequality (1.14) becomes inequalities (1.10) and (1.11), respectively.

2 Lemmas

In order to prove Theorems 1.1 and 1.2 we need two lemmas, which we present in this section.

Lemma 2.1. Let $u \in [0, 1]$, $s \geq 1$ and

$$f_{u,s}(x) = \frac{s}{2} \log(1 - ux^2) - \log\left(\frac{\pi}{2\mathcal{K}(x)}\right). \quad (2.1)$$

Then $f_{u,s} > 0$ for all $x \in (0, 1)$ if and only if $2su \leq 1$.

Proof. From (1.4) and (2.1) one has

$$f'_{u,s}(x) = -\frac{usx}{1 - ux^2} + \frac{\mathcal{E}(x) - (1 - x^2)\mathcal{K}(x)}{x(1 - x^2)\mathcal{K}(x)} = \frac{F_{u,s}(x)}{x(1 - x^2)(1 - ux^2)\mathcal{K}(x)}, \quad (2.2)$$

where

$$F_{u,s}(x) = -sux^2(1 - x^2)\mathcal{K}(x) + (1 - ux^2)[\mathcal{E}(x) - (1 - x^2)\mathcal{K}(x)]. \quad (2.3)$$

It follows from (1.1)-(1.3) and (2.3) together with elaborated computations that

$$\begin{aligned} & \mathcal{E}(x) - (1 - x^2)\mathcal{K}(x) \\ &= \frac{\pi}{2} \left[\sum_{n=0}^{\infty} \frac{(-1/2, n)(1/2, n)}{(n!)^2} x^{2n} - (1 - x^2) \sum_{n=0}^{\infty} \frac{(1/2, n)^2}{(n!)^2} x^{2n} \right] \\ &= \frac{\pi}{2} \sum_{n=0}^{\infty} \frac{(1/2, n)^2}{2n!(n+1)!} x^{2n+2}, \end{aligned}$$

$$\begin{aligned}
\frac{2}{\pi} F_{u,s}(x) &= -su x^2(1-x^2) \sum_{n=0}^{\infty} \frac{(1/2, n)^2}{(n!)^2} x^{2n} + (1-ux^2) \sum_{n=0}^{\infty} \frac{(1/2, n)^2}{2n!(n+1)!} x^{2n+2} \\
&= -su \sum_{n=0}^{\infty} \frac{(1/2, n)^2}{(n!)^2} x^{2n+2} + su \sum_{n=0}^{\infty} \frac{(1/2, n)^2}{(n!)^2} x^{2n+4} \\
&\quad + \sum_{n=0}^{\infty} \frac{(1/2, n)^2}{2n!(n+1)!} x^{2n+2} - u \sum_{n=0}^{\infty} \frac{(1/2, n)^2}{2n!(n+1)!} x^{2n+4} \\
&= -su x^2 - su \sum_{n=0}^{\infty} \frac{(1/2, n+1)^2}{[(n+1)!]^2} x^{2n+4} + su \sum_{n=0}^{\infty} \frac{(1/2, n)^2}{(n!)^2} x^{2n+4} \\
&\quad + \frac{x^2}{2} + \sum_{n=0}^{\infty} \frac{(1/2, n+1)^2}{2(n+1)!(n+2)!} x^{2n+4} - u \sum_{n=0}^{\infty} \frac{(1/2, n)^2}{2n!(n+1)!} x^{2n+4} \\
&= x^2 \left[\frac{1}{2} - su + \sum_{n=0}^{\infty} \frac{(1/2, n)^2 A_n}{2(n+1)!(n+2)!} x^{2n+2} \right], \tag{2.4}
\end{aligned}$$

where

$$A_n = su(n+2)(2n + \frac{3}{2}) + (n + \frac{1}{2})^2 - u(n+1)(n+2) > 0. \tag{2.5}$$

We divide the proof into two cases.

Case 1.1. $2su \leq 1$. Then (2.2)-(2.5) lead to conclusion that $f_{u,s}(x)$ is strictly increasing in $(0, 1)$. Therefore, $f_{u,s}(x) > f_{u,s}(0^+) = 0$ for all $x \in (0, 1)$ follows from (1.2) and (2.1) together with the monotonicity of $f_{u,s}(x)$ in $(0, 1)$.

Case 1.2. $2su > 1$. Then (2.2)-(2.4) lead to conclusion that there exists $\delta_1 \in (0, 1)$ such that $f_{u,s}(x)$ is strictly decreasing in $(0, \delta_1)$. Therefore, $f_{u,s}(x) < f_{u,s}(0^+) = 0$ for all $x \in (0, \delta_1)$ follows from (1.2) and (2.1) together with the monotonicity of $f_{u,s}(x)$ in $(0, \delta_1)$.

Lemma 2.2. Let $u \in [0, 1]$, $s \geq 1$, $\operatorname{arctanh}(x) = \frac{1}{2} \log \left(\frac{1+x}{1-x} \right)$ be the inverse hyperbolic tangent function, and

$$g_{u,s}(x) = \frac{s}{2} \log(1-ux^2) + \log \left(\frac{\operatorname{arctanh}(x)}{x} \right). \tag{2.6}$$

Then $g_{u,s}(x) > 0$ for all $x \in (0, 1)$ if and only if $3su \leq 2$.

Proof. From (2.6) one has

$$g'_{u,s}(x) = -\frac{sux}{1-ux^2} + \frac{x - (1-x^2)\operatorname{arctanh}(x)}{x(1-x^2)\operatorname{arctanh}(x)} = \frac{G_{u,s}(x)}{x(1-x^2)(1-ux^2)\operatorname{arctanh}(x)}, \tag{2.7}$$

where

$$G_{u,s}(x) = -sux^2(1-x^2)\operatorname{arctanh}(x) + (1-ux^2)[x - (1-x^2)\operatorname{arctanh}(x)]. \quad (2.8)$$

Making use of series expansion and (2.8) we have

$$\begin{aligned} G_{u,s}(x) &= -sux^2(1-x^2) \sum_{n=0}^{\infty} \frac{x^{2n+1}}{2n+1} + (1-ux^2) \left[x - (1-x^2) \sum_{n=0}^{\infty} \frac{x^{2n+1}}{2n+1} \right] \\ &= -su \sum_{n=0}^{\infty} \frac{x^{2n+3}}{2n+1} + su \sum_{n=0}^{\infty} \frac{x^{2n+5}}{2n+1} + (1-ux^2) \sum_{n=0}^{\infty} \frac{2x^{2n+3}}{(2n+1)(2n+3)} \\ &= x^3 \left[\frac{2}{3} - su + \sum_{n=0}^{\infty} \frac{B_n x^{2n+2}}{(2n+1)(2n+3)(2n+5)} \right], \end{aligned} \quad (2.9)$$

where

$$B_n = 2u(s-1)(2n+5) + 2(2n+1) > 0. \quad (2.10)$$

We divide the proof into two cases.

Case 1.1. $3su \leq 2$. Then (2.7)-(2.10) lead to conclusion that $g_{u,s}(x)$ is strictly increasing in $(0, 1)$. Therefore, $g_{u,s}(x) > g_{u,s}(0^+) = 0$ for all $x \in (0, 1)$ follows from (2.6) together with the monotonicity of $g_{u,s}(x)$ in $(0, 1)$.

Case 1.2. $3su > 2$. Then (2.7)-(2.9) lead to conclusion that there exists $\delta_2 \in (0, 1)$ such that $g_{u,s}(x)$ is strictly decreasing in $(0, \delta_2)$. Therefore, $g_{u,s}(x) < g_{u,s}(0^+) = 0$ for all $x \in (0, \delta_2)$ follows from (2.6) and the monotonicity of $g_{u,s}(x)$ in $(0, \delta_2)$.

3 Proof of Theorems 1.1 and 1.2

Proof of Theorem 1.1. Since both $Q_{t,s}(a, b)$ and $AG(a, b)$ are symmetric and homogeneous of degree 1. Without loss of generality, we assume that $a > b$. Let $x = (a-b)/(a+b) \in (0, 1)$. Then from (1.5) and (1.7) together with $b/a = (1-x)/(1+x)$ we have

$$\begin{aligned} \frac{AG(a, b)}{A(a, b)} &= \frac{AG(1, b/a)}{A(1, b/a)} = \frac{\pi}{\mathcal{K} \sqrt{1 - (b/a)^2 (1 + b/a)}} \\ &= \frac{\pi(1+x)}{2\mathcal{K}(2\sqrt{x}/(1+x))} = \frac{\pi}{2\mathcal{K}(x)}. \end{aligned} \quad (3.1)$$

It follow from (1.12) and (3.1) that

$$\begin{aligned}\log\left(\frac{Q_{t,s}(a,b)}{AG(a,b)}\right) &= \log\left(\frac{Q_{t,s}(a,b)}{A(a,b)}\right) - \log\left(\frac{AG(a,b)}{A(a,b)}\right) \\ &= \frac{s}{2} \log[1 - (1-2t)^2 x^2] - \log\left[\frac{\pi}{2\mathcal{K}(x)}\right].\end{aligned}\quad (3.2)$$

Therefore, Theorem 1.1 follows from Lemma 2.1 and (3.2).

Proof of Theorem 1.2. Since both $Q_{t,s}(a,b)$ and $L(a,b)$ are symmetric and homogeneous of degree 1. Without loss of generality, we assume that $a > b$. Let $x = (a-b)/(a+b) \in (0,1)$. Then (1.12) leads to

$$\begin{aligned}\log\left(\frac{Q_{t,s}(a,b)}{L(a,b)}\right) &= \log\left(\frac{Q_{t,s}(a,b)}{A(a,b)}\right) - \log\left(\frac{L(a,b)}{A(a,b)}\right) \\ &= \frac{s}{2} \log[1 - (1-2t)^2 x^2] + \log\left(\frac{\operatorname{arctanh}(x)}{x}\right).\end{aligned}\quad (3.3)$$

Therefore, Theorem 1.2 follows from Lemma 2.2 and (3.3).

Acknowledgement: This work was supported by the Natural Science Foundation of China (Grant Nos. 11071059, 11071069, 11171307), and the Innovation Team Foundation of the Department of Education of Zhejiang Province (Grant no. T200924).

References

- [1] F. Bowman, Introduction to Elliptic Functions with Application, Dover Publications, New York, 1961.
- [2] P. F. Byrd and M. D. Friedman, Handbook of Elliptic Integrals for Engineers and Scientists. Springer-Verlag, New York, 1971.
- [3] G. D. Anderson, M. K. Vamanamurthy and M. Vuorinen, Conformal Invariants, Inequalities, and Quasiconformal Maps, John Wiley & Sons, New York, 1997.
- [4] W. Fechner, On some functional inequalities related to the logarithmic mean, Acta Math. Hungar., 2010, **128**(1-2): 36-45.

- [5] H. Kosaki, Arithmetic-geometric mean and related inequalities for operators, *J. Funct. Anal.*, 1998, **156**(2): 429-451.
- [6] J. Sándor, On certain inequalities for means II, *J. Math. Anal. Appl.*, 1996, **199**(2): 629-635.
- [7] J. Sándor, On certain inequalities for means, *J. Math. Anal. Appl.*, 1995, **189**(2): 602-606.
- [8] L. G. Lucht, On the arithmetic-geometric mean inequality, *Amer. Math. Monthly*, 1995, **102**(8): 739-740.
- [9] E. Neuman, The weighted logarithmic mean, *J. Math. Anal. Appl.*, 1994, **188**(3): 885-900.
- [10] J. Sándor, On the identric and logarithmic means, *Aequationes Math.*, 1990, **40**(2-3): 261-270.
- [11] T. P. Lin, The power mean and the logarithmic mean, *Amer. Math. Monthly*, 1974, **81**: 879-883.
- [12] B. C. Carlson, The logarithmic mean, *Amer. Math. Monthly*, 1972, **79**: 615-618.
- [13] B. C. Carlson, Algorithms involving arithmetic and geometric means, *Amer. Math. Monthly*, 1971, **78**: 496-505.
- [14] B. C. Carlson and M. Vuorinen, Inequalities of the AGM and the logarithmic mean, *SIAM Review*, 1991, **33**(4): 655-655.
- [15] P. Bracken, An arithmetic-geometric mean inequality, *Expo. Math.*, 2001, **19**(3): 273-279.
- [16] M. K. Vamanamurthy and M. Vuorinen, Inequalities for means, *J. Math. Anal. Appl.*, 1994, **183**(1): 155-166.
- [17] Y.-M. Chu and M.-K. Wang, Optimal inequalities between harmonic, geometric, logarithmic, and arithmetic-geometric means, *J. Appl. Math.*, 2011, Article ID 618929, 9 pages.
- [18] Y.-M. Chu, M.-K. Wang and Z.-K. Wang, Best possible inequalities among harmonic, geometric, logarithmic and Seiffert means, *Math. Inequal. Appl.*, 2012, **15**(2): 415-422.